

Sample Date  
3/16/2011



**BUD KILL**

Variety	Map Location #	Stage	Control	26	23	21	19
Mid Valley Bing	12	2+	0%	5%	12%	42%	81%
			0 137	6 126	16 135	44 106	122 151
Mid Valley Rainier	11	2	0%	2%	6%	6%	50%
			0 56	3 143	8 132	4 65	92 184
Mid Valley Chelan	11	1	0%	0%	1%	0%	14%
			0 136	0 141	2 162	0 200	21 153
Mid Valley Duke	12		9%	7%	0%	8%	15%
			13 144	7 106	0 103	13 155	18 119
Wapato Supreme	14	3	10%	12%	18%	54%	62%
			5 50	3 26	6 34	14 26	13 21
Wapato Rich Lady	14	3	24%	57%	51%	44%	32%
			9 38	28 49	22 43	18 41	13 41
Mid Valley Gala	12	1	0%	6%	2%	15%	75%
			0 52	2 36	1 42	9 59	21 28
Mid Valley Bartletts	12		0%	0%	0%	10%	8%
			0 57	0 64	0 58	6 61	4 48
Upper Valley Sweetheart	17	1	0%	0%	9%	4%	25%
			0 66	0 153	18 203	4 107	49 196

The provided information shows percent bud damage at various temperatures in the bud wood selected. This information may not indicate the exact temperature sensitivity of the buds in your orchard. Location, variety, tree health, tree age, rootstock, cultural practices, dew point, and duration of cold can all have an influence on critical temperatures.

\*\*Another historical reference to consider when making your critical temperature decision would be the

**Critical Temperature for Blossom Buds.** This can be accessed at: <http://treefruit.yakima.wsu.edu> under **Tree Fruit Bud Charts.**

Sample Date  
3/16/2011



**BUD KILL**

Variety	Map Location #	Stage	Control	26	23	21	19
<b>Pasco Snow Giant</b>	<b>6</b>	<b>3</b>	<b>13%</b>	<b>42%</b>	<b>70%</b>	<b>44%</b>	<b>28%</b>
			2 15	5 12	14 20	16 36	7 25
<b>Pasco Sweet Dream</b>	<b>6</b>	<b>3</b>	<b>64%</b>	<b>85%</b>	<b>67%</b>	<b>92%</b>	<b>73%</b>
			9 14	17 20	14 21	22 24	11 15
<b>Pasco Summer Sweet</b>	<b>6</b>	<b>3</b>	<b>36%</b>	<b>16%</b>	<b>47%</b>	<b>100%</b>	<b>88%</b>
			5 14	3 19	8 17	10 10	22 25
<b>Pasco Summer Blush</b>	<b>6</b>	<b>3</b>	<b>17%</b>	<b>17%</b>	<b>63%</b>	<b>80%</b>	<b>93%</b>
			3 18	2 12	10 16	12 15	14 15
<b>Outlook Bing</b>	<b>10</b>	<b>2</b>	<b>0%</b>	<b>1%</b>	<b>46%</b>	<b>39%</b>	<b>29%</b>
			0 45	1 79	28 61	35 90	30 103
<b>Richland Chelan</b>	<b>7</b>	<b>2</b>	<b>0%</b>	<b>1%</b>	<b>13%</b>	<b>19%</b>	<b>33%</b>
			0 106	1 144	15 114	22 116	42 128
<b>Wallula Chelan</b>	<b>1</b>	<b>2</b>	<b>0%</b>	<b>16%</b>	<b>24%</b>	<b>39%</b>	<b>93%</b>
			0 59	16 100	25 103	56 143	50 54
<b>Wallula Pink lady</b>	<b>1</b>	<b>2+</b>	<b>5%</b>	<b>19%</b>	<b>38%</b>	<b>44%</b>	<b>100%</b>
			2 44	13 67	18 48	18 41	34 34
<b>Wallula Gala</b>	<b>1</b>	<b>2</b>	<b>0%</b>	<b>38%</b>	<b>78%</b>	<b>89%</b>	<b>67%</b>
			0 54	25 65	45 58	56 63	54 81
<b>Snake River Chelan Ice Harbor</b>	<b>4</b>	<b>1+</b>	<b>0%</b>	<b>3%</b>	<b>1%</b>	<b>57%</b>	<b>80%</b>
			0 56	5 177	2 282	54 95	113 141
<b>Snake River Bing Ice Harbor</b>	<b>4</b>	<b>2</b>	<b>0%</b>	<b>0%</b>	<b>2%</b>	<b>18%</b>	<b>46%</b>
			0 124	0 83	2 99	16 88	26 57

The provided information shows percent bud damage at various temperatures in the bud wood selected. This information may not indicate the exact temperature sensitivity of the buds in your orchard. Location, variety, tree health, tree age, rootstock, cultural practices, dew point, and duration of cold can all have an influence on critical temperatures.

\*\*Another historical reference to consider when making your critical temperature decision would be the

**Critical Temperature for Blossom Buds.** This can be accessed at: <http://treefruit.yakima.wsu.edu> under **Tree Fruit Bud Charts**.



Sample Date  
3/16/2011

**BUD KILL**

Variety	Map Location #	Stage	Control	26	23	21	19
Quincy Sweethearts		Dormant	2%	0%	10%	2%	2%
			1 56	0 100	6 62	2 106	4 191
Quincy Cotts			6%	7%	18%	28%	73%
			2 31	4 57	7 39	19 68	22 30
Prosser Bing	9	1+	0%	3%	3%	31%	18%
			0 213	6 197	7 201	49 157	34 192
Prosser Chelans	9	1+	0%	0%	1%	7%	42%
			0 122	0 166	3 209	9 126	57 135
Benton City Bing	8	2	1%	2%	24%	14%	72%
			1 147	4 195	11 46	13 90	61 85
Benton City Chelan	8	1+	0%	0%	15%	24%	68%
			0 130	0 95	28 186	40 164	82 120
Kennewick Chelans	5	3	0%	9%	36%	48%	27%
			0 85	10 112	16 44	65 135	37 138
Kennewick Bing	5	2	0%	10%	29%	43%	78%
			0 67	9 89	24 84	40 94	46 59
Burbank Chelan	2	3	1%	57%	62%	71%	69%
			2 177	72 127	73 118	141 198	164 238
Mattawa Chelans	21	2	24%	13%	29%	67%	32%
			26 109	18 141	37 127	163 244	58 179
Mattawa Benton	21	1	3%	11%	22%	18%	15%
			8 239	24 217	48 222	29 159	29 194
Finley Early Robins	5	3	4%	11%	34%	86%	97%
			4 106	11 97	31 91	76 88	98 101

The provided information shows percent bud damage at various temperatures in the bud wood selected. This information may not indicate the exact temperature sensitivity of the buds in your orchard. Location, variety, tree health, tree age, rootstock, cultural practices, dew point, and duration of cold can all have an influence on critical temperatures.

\*\*Another historical reference to consider when making your critical temperature decision would be the

**Critical Temperature for Blossom Buds.** This can be accessed at: <http://treefruit.yakima.wsu.edu> under **Tree Fruit Bud Charts.**