



FOLI-GRO[®]

LIQUID FOLIAR FERTILIZER



3-18-18

3-18-18

GUARANTEED ANALYSIS

Total Nitrogen (N)	3.0%
0.50% Ammoniacal Nitrogen	
2.50% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	18.0%
Soluble Potash (K ₂ O)	18.0%
Derived from: Ammonia, Urea, Phosphoric Acid, Potassium Hydroxide.	

KEEP OUT OF REACH OF CHILDREN

CAUTION:

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. In case of contact, immediately flush skin or eyes with water for at least 15 minutes.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-we.com/>

FIRST AID

In all cases, call a poison control center or doctor for further treatment advice.

IF SWALLOWED, call a poison control center or doctor immediately. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **IF ON SKIN**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration. **IF IN EYES**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Store in a safe manner. Store in original container only and keep tightly sealed when not in use. Dispose of unused product and empty containers in accordance with Federal, State and local regulations.

GENERAL INFORMATION

FOLI-GRO[®] is manufactured specifically for use as a liquid foliar or soil applied fertilizer. Its use is suggested as part of a Total Nutrition System[®] program to enhance yields and quality.

Under adverse environmental conditions (water stress, cold soils, salts, etc.) when used as directed FOLI-GRO[®] may supply otherwise unavailable plant nutrients. Research has shown that foliarly applied nutrients are absorbed more efficiently by plants than are those supplied in the soil under the above conditions.

FOLI-GRO[®] is uniquely formulated to supply nutrients in a readily available form for the plant to obtain a quick response to needed elements. A soil and tissue testing program may be helpful to monitor and maintain optimum plant growth and quality. Nutrient translocation inside the plant is generally more rapid when applied foliarly.

3-18-18 is a highly soluble, low salt index formulation developed to supplement standard soil fertility practice. Provides an additional source of nitrogen, phosphorous and potassium nutrients during the growing season.

3-18-18 is designed to be used in concert with soil and tissue tests and professional recommendations to manage the nutrient levels and nutrient balance within the crop.

3-18-18 is in the form readily absorbed by plant tissue. In this form, nitrogen, phosphorous and potassium can be applied to the growing plant.

SUGGESTIONS FOR USE

FIELD CROPS (Foliar Application)

Alfalfa: One to three gallons/acre immediately after each cutting. Two to three gallons per acre after the final cutting of the season.

Beans, Peas: One gallon/acre at first true leaf stage, one to two gallons/acre prebloom; and one to two gallons/acre at pod fill.

Corn: One to three gallons/acre at 8-12 inch and 24-30 inch height.

Potatoes: One to three gallons/acre three to four times during season. The first at early emergence, three to four weeks later and prebloom.

SUGGESTIONS FOR USE (cont.)

Safflower: One to three gallons/acre when four to eight inches tall and two to three weeks prior to bloom.

Soybeans: One to three gallons/acre, three times each season starting at first full leaf. Last application should be at early pod fill.

Wheat, Barley, Oats, and Rice: One to two gallons/acre at early tillering and again at early boot stage.

VEGETABLE AND HORTICULTURAL CROPS (Foliar Application)

Asparagus: One to three gallons/acre at full fern and three to four gallons/acre two weeks before fern turns yellow in the fall.

Bush Berries: One to three gallons/acre three times each season. Starting at first full leaf, then post-bloom, and at berry sizing time.

Celery: One to three gallons/acre immediately after transplanting or thinning, three to four weeks after first application and three to four weeks prior to harvest.

Cole Crops: (Cabbage, Cauliflower, Broccoli, Brussels Sprouts) One to three gallons/acre immediately after transplanting or thinning, three to four weeks after first application and three to four weeks prior to harvest.

Letuce: One to three gallons/acre two to three times each season. Starting at second true leaf after transplanting or thinning, at folding, with the last application three weeks prior to harvest.

Melons, Cucumbers, Squash: One to three gallons/acre three times per year. Starting at first new growth after transplanting or thinning. Then at first blossom, and at fruit set/fruit size.

Onions, Garlic: One to three gallons/acre two to three times each season. Starting when first leaf is 3 inches, then at midseason, and then two and three weeks prior to harvest.

Peppers: One to three gallons/acre two to three times each season. Starting at first new growth after transplant or thinning. Then at first blossom, and at fruit set/fruit size.

Spinach: One to three gallons/acre two to three times each season. Starting at first true leaf, then midseason and two to three weeks prior to harvest.

Strawberries: One to three gallons/acre at early fruit set and after each picking.

Sweet Potatoes: Use planting drench of three gallons/acre/100 gallons of water, then foliar one to three gallons/acre three weeks after planting, midseason, and two to three weeks prior to harvest.

Tomatoes: One to three gallons/acre four times each season. Starting at first true leaf then at early bloom, fruit set, and early fruit size.

TRANSPLANT SOLUTIONS AND DRENCHES

Mix thoroughly one to three gallons in not less than 100 gallons of water and drench roots. For vegetables drench entire plant. Plant immediately after drenching. Do not allow plants to dry or wilt. Total amount of 3-18-18 used should not exceed 3 gallons per acre regardless of amounts of water used in transplanting.

TREE CROPS (Foliar Application)

Apples, Nectarines, Peaches, Pears, and Plums: One to three gallons/acre three to four times each season. Starting at three-quarter leaf, then at early fruit size, fruit midsize and three to four weeks prior to harvest.

Cherries and Apricots: One to three gallons/acre three to four times each season. Starting at pink bud, first full leaf, then at fruit midsize and post-harvest.

SUGGESTIONS FOR USE (cont.)

Walnuts, Almonds, Pistachios, Filberts and Pecans: One to four gallons/acre three times each season starting at first full leaf, early nut development, and three to four weeks prior to hull split.

TURF GRASSES (Foliar Application)

One gallon per 5000 square feet applied with sufficient water to insure uniform distribution. May be applied every 30 to 60 days during growing season. Supplemental nitrogen should be added as recommended by your supplier and/or Pest Control Advisor.

VINE CROPS (Foliar Application)

Grapes: One to three gallons/acre at first full leaf, then two to three gallons/acre at post bloom, berry size and three to four weeks prior to harvest.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS;" AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation uses contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS Logo[®], IDEAS TO GROW WITH[®], FOLI-GRO[®] and TOTAL NUTRITION SYSTEM[®] are registered trademarks of Wilbur-Ellis Company.

Manufactured by Wilbur-Ellis Company
7 E. Washington Ave. • Yakima, WA 98903

P-12707

NET CONTENTS:

Lot #

GALLONS (

WEIGHT 11.80 POUNDS/GAL.

Liters)