



WILBUR-ELLIS®

VENGEANCE® PLUS

SELECTIVE HERBICIDE

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES, SOD FARMS, NON-TURF AREAS, INDUSTRIAL VEGETATION MANAGEMENT AND NON-CROPLAND USE

CONTAINS MCPA, TRICLOPYR AND DICHLORPROP-p

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*.....	46.87%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**.....	10.68%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy) propionic Acid***‡.....	9.12%

OTHER INGREDIENTS:33.33%
100%

By Isomer specific AOAC Method Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid 38.27%, 3.72 lbs./gal.

**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid 7.65%, 0.75 lbs./gal.

***(+)-R-2-(2,4-Dichlorophenoxy) propionic Acid 7.65%, 0.75 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p

For Use By Turf Maintenance Personnel, Landscaping or Commercial Applicators Only

EPA REG. NO. 228-411-2935

EPA EST. NO. 228-IL-1

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

For Chemical Spill, Leak, Fire or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO

Corrosive. Causes irreversible eye damage. May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

NON-WPS TURF USE: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general only agricultural plant uses are covered by the WPS - must wear: coveralls over short-sleeve shirt and short pants, socks, chemical-resistant footwear and chemical-resistant gloves made of any waterproof material, eye protection (goggles, face shield, or safety glasses) when using this product. It is recommended that safety glasses include front, brow, and temple protection.

WPS SOD FARM USES: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) – in general, agricultural plant uses are covered - must wear: coveralls over short-sleeve shirt and short pants, socks, chemical-resistant footwear and chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride and protective eyewear. Some materials that are chemical-resistant to this product are made out of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance selection chart.

Mixers, loaders, applicators, flaggers and other handlers must wear:

- Coverall over short-sleeved shirt and short pants
- Protective eyewear (goggles or face shield)
- Chemical-resistant gloves made of any any water-proof material
- Chemical-resistant footwear and socks
- When mixing and loading wear chemical-resistant apron
- For overhead exposure wear chemical-resistant headgear
- When cleaning equipment add a chemical-resistant apron

Engineering Controls Statements: If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	

NET CONTENTS: 2.5 GALLONS

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants and may be hazardous to aquatic organisms in water adjacent to treated areas. MCPA may be toxic to fish, aquatic invertebrates and aquatic plants. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollution Discharge Elimination (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Aerial application of this product is prohibited.

Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs) and similar turf areas. *Excluding Bentgrass Tees.

For residential turf and golf course uses, do not apply more than 1.5 lb of MCPA (3 pints) per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, grapes, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede or St. Augustine grasses unless turf injury can be tolerated. The suitable use of this product on nonrecommended turf species may be determined by treating a small area at any rate per acre which does not exceed 3 pints per acre. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays

WEEDS CONTROLLED BY VENGEANCE® PLUS SELECTIVE HERBICIDE

This product is designed to control many broadleaf weeds such as:

Alder	Bracted plantain	Catsear
Annual yellow sweet clover	Brassbuttons	Catnip
Artichoke	Bristly oxtongue	Chickweed
Aster	Broadleaf dock	Chicory
Austrian Fieldcress	Broadleaf plantain	Cinquefoil
Bedstraw	Broomweed	Clover
Beggartick	Buckhorn	Cockle
Biden	Buckhorn plantain	Cocklebur
Bindweed	Bulbous buttercup	Coffeebean
Bird vetch	Bull nettle	Coffeeweed
Bitterweed	Bull thistle	Common chickweed
Bitter wintercress	Burdock	Common mullein
Black-eyed Susan	Burning nettle	Common sowthistle
Black medic	Bur ragweed	Corn chamomile
Black mustard	Burweed	Creeping jenny
Blackseed plantain	Buttercup	Crimson clover
Blessed thistle	Canada thistle	Croton
Blue lettuce	Carolina geranium	Cudweed
Blue vervain	Carpetweed	Curly dock
Box elder	Catchweed	Curly indigo
	bedstraw	Dandelion

Continued:

Dead nettle	Narrowleaf vetch	Spurge
Dock	Nettle	Spurweed
Dogbane	Orange hawkweed	Stinging nettle
Dogfennel	Oxalis	Stinkweed
Elderberry	Oxeye daisy	Stitchwort
English daisy	Parsley-piert	Strawberry clover
Fall dandelion	Parsnip	Sumac
False dandelion	Pearlwort	Sunflower
False flax	Pennycress	Sweet clover
False sunflower	Pennywort	Tall nettle
Fiddleneck	Peppergrass	Tall vervain
Field bindweed	Pepperweed	Tansy mustard
Field pansy	Pigweed	Tansy ragwort
Flea bane (daisy)	Pineywoods bedstraw	Tanweed
Flixweed	Plains coreopsis	Tanweed
Florida betony	Plantain	Thistle
Florida pusley	Poison hemlock	Tick trefoil
Frenchweed	Poison ivy	Toadflax
Galinsoga	Poison oak	Trailing crownvetch
Garlic mustard	Pokeweed	Tumble mustard
Goathead	Poorjoe	Tumble pigweed
Goatsbeard	Povertyweed	Tumbleweed
Goldenrod	Prickly lettuce	Velvet leaf
Ground ivy	Prickly sida	Venice mallow
Gumweed	Primrose	Veronica
Hairy bittercress	Prostrate knotweed	Vervain
Hairy fleabane	Prostrate pigweed	Vetch
Hawkweed	Prostrate spurge	Virginia buttonweed
Healall	Prostrate vervain	Virginia creeper
Heartleaf drymary	Puncture vine	Virginia pepperweed
Hedge bindweed	Purslane	Wavyleaf bullthistle
Hedge mustard	Ragweed	Western clematis
Hemp	Red clover	Western salsify
Henbit	Redroot pigweed	White clover
Hoary cress	Red sorrel	White mustard
Hoary plantain	Remstem filaree	Wild mustard
Hoary vervain	Rough cinquefoil	Wild aster
Honeysuckle	Rough fleabane	Wild buckwheat
Hop clover	Roundleafed marigold	Wild carrot
Horsenettle	Rush	Wild four-o'clock
Horsetail	Russian pigweed	Wild garlic
Indiana mallow	Russian thistle	Wild geranium
Ironweed	St. Johnswort	Wild lettuce
Jewelweed	Scarlet pimpernel	Wild marigold
Jimsonweed	Scotch thistle	Wild onion
Kochia	Sheep sorrel	Wild parsnip
Knawel	Shepherdspurse	Wild radish
Knotweed	Slender plantain	Wild rape
Lambsquarter	Smallflower galinsoga	Wild strawberry
Lespedeza	Smartweed	Wild sweet potato
Locoweed	Smooth dock	Wild vetch
Lupine	Smooth pigweed	Willow
Mallow	Sneezeweed	Witchweed
Marshelder	Southern wild rose	Woolly morningglory
Matchweed	Sowthistle	Woodsorrel
Mexicanweed	Spanishneedle	Woolly croton
Milk vetch	Spatdock	Woolly plantain
Milkweed bloodflower	Speedwell	Wormseed
Mugwort	Spiny amaranth	Yarrow
Morningglory	Spiny cocklebur	Yellow rocket
Mouseear chickweed	Spotted catsear	Yellowflower pepperweed
Musk thistle	Spotted knapweed	and other broadleaf weeds
Mustard	Spotted spurge	
Narrowleaf plantain		

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf grasses planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use this product at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

NOTE: For all grasses: (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

SOD FARMS

For sod farms, do not apply more than 1.5 lb of MCPA (3 pints) per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

This product is intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

Timing

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to Floratam St. Augustinegrass.

The application of this herbicide to grass seedlings is recommended after the second mowing.

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 or 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant Bermudagrass, dormant Zoysiagrass and dormant Bahiagrass are suggested.

Rates for Sod Farms

For Bluegrass, Ryegrass and Fescue, apply 1-1/2 to 3 pints per acre. For Creeping Bentgrass use .9 pints per acre. For Common and Hybrid Bermuda, Bahia, Zoysia and St. Augustine grass apply 1 to 2 pints per acre. Spray volume for Sod Farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

NON-TURF AREAS

For non-cropland Rights-of-Way use, do not apply more than 1.5 lb ae/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

For spot treatment noncropland Rights-of-Way use, do not apply more than 3.0 lb ae/acre per year.

Roadsides, Rights-of-Way, and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1.6 to 4.8 pints of this product in sufficient volume of water to ensure thorough coverage. Apply any time after the plants come into full leaf stage. Best results are obtained when weeds are actively growing. For (spot) applications with small tank sprayers, apply at the rate of 2.25 ounces of this product per gallon of water and spray foliage to wet.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 4.8 to 7.2 pints of this product to 100 gallons of water applying 200 to 600 gallons of spray mixture per acre depending on the height and thickness of the brush. Mix thoroughly before spraying.

COMPATIBILITY

A compatibility test is recommended prior to mixing with liquid fertilizers and liquid iron. All regulations, either state or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

INDUSTRIAL VEGETATION MANAGEMENT

Water soluble industrial herbicide for the control of brush and broadleaf weeds on Rights-of-Way, Railroads, Pipelines, Brush, Fence Rows, Non-Irrigation Ditchbanks, Wasteland and similar noncropland areas.

NONCROPLAND USE: (Roadsides, fence rows, railroads, brush control or forest site preparation and similar areas.) For top growth control of field bindweed, Canada thistle, perennial sowthistle, Russian knapweed and leafy spurge, apply 1.6 to 4.8 pints of this product per acre.

Continued:

FOR SPOT TREATMENT: Apply 1 teaspoonful of this product per 100 to 275 square feet. Apply as a foliar spray to actively growing plants.

BRUSH, BROADLEAF, INVASIVE AND NOXIOUS WEED CONTROL

SPECIES CONTROLLED

When used as directed, this product will control or suppress many herbaceous broadleaf weeds (annual, biennial, and perennial) as well as suppress many unwanted woody plant and vine species. Species controlled include:

ANNUALS

Buckwheat	Knawel	Purslane
Carpeweed	Kochia	Ragweed
Chickweed	Lambsquarter	Russian Thistle
Clover	Morningglory	Smartweed
Cocklebur	Mustard	Velvetleaf
Daisy, English	Pigweed	
Henbit	Puncturevine	

Biennials

Ragwort, Tansy Thistle, Musk

PERENNIALS

Bindweed, Field	Dogfennel	Sorrel, Sheep
Carrot, Wild	Knapweed, Russian	Spurge, Leafy
(Queen Anne's Lace)	Milkweed	Thistle, Canada
Dock, Curly	Ragweed, Perennial	Toadflax, Dalmation

WOODY BRUSH AND VINES*

Alder	Gum	Poplar	Spruce
Ash	Hawthorn	Puncturevine	Sumac
Aspen	Hemlock	Raspberry	Sycamore
Basswood	Honeysuckle	Redcedar, Eastern	Trumpet creeper
Beech	Ivy, Poison	Redvine	Waxmyrtle
Birch	Kudzu	Rose, Multiflora	Willow
Blackberry	Locust	Sagebrush	Witchhazel
Cherry	Maple	Sassafras	Yaupon
Creeper, Virginia	Oak	Schinus (Florida Holly,	
Creosotebush	Olive, Russian	Brazil Peppertree,	
Cucumbertree	Persimmon	Christmas-berry)	
Dogwood	Pine	Serviceberry	
Elderberry	Plum, Wild	Snowberry	
Elm			

*Suppression

APPLICATION DIRECTIONS

Regardless of the species to be controlled, a sufficient volume of mixture should be used to allow for thorough spray coverage. Make applications when weeds and brush are actively growing. The addition of a Chemical Producers and Distributors Association's (CPDA) certified adjuvant can increase control. A CPDA certified drift control agent may also be used. Biennials are best controlled when treated in the rosette stage. Regrowth may occur on resistant species. Retreatments may be made as needed; however, do not exceed a total of 4.8 pints of this product per treated acre during a growing season. For control of woody species, a tank mix may be necessary. Tank mix recommendations are for use only in states where the tank mix product and application site are registered. Read and follow all label directions.

HERBACEOUS BROADLEAF WEED CONTROL

Apply 1.6 to 4.8 pints of this product in sufficient volume of water to allow for thorough coverage of weeds. Do not apply more than 4.8 pints of product per treated acre.

WOODY BRUSH AND VINE CONTROL

High Volume Foliar Spot Applications: Mix 4.8 to 7.2 pints of this product in sufficient volume of water to allow for thorough coverage. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 7.2 pints of product per treated acre. Direct the spray to treat all foliage, stems, and root collars to wet.

WOODY BRUSH AND VINE CONTROL/SUPPRESSION

Broadcast Applications with Ground Equipment: Apply 4.8 to 7.2 pints of this product in sufficient volume of water to allow for thorough coverage per treated acre. Do not apply more than 7.2 pints of product per treated acre. Spray all foliage, stems, and root collars to wet. For optimum control of woody species, a tank mix may be necessary.

TANK MIX TREATMENTS

TANK MIX RECOMMENDATIONS ARE FOR USE ONLY IN STATES WHERE THE TANK MIX PRODUCT AND APPLICATION SITE ARE REGISTERED. READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, AND OTHER RESTRICTIONS. For broader spectrum control, this product may be tank mixed with other herbicides registered for noncropland uses (e.g. railroad, highway, pipeline, forest brush control, etc.), including but not limited to the following herbicides, in accordance with the more restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Add water to the spray tank prior to the addition of the tank mix products. Do not premix concentrates.

HERBICIDE

Asulam (Asulox®)	Metsulfuron Methyl (Patriot®)**
Bromacil (Hyvar®)	MSMA
Chlorsulfuron (Telar®)	Picloram (Tordon®)*
Dicamba (Diablo™)	Simazine (Princep®)*
Diuron (Karmex®)	Sulfometuron methyl (Spyder™)
Fosamine ammonium (Krenite®)	Tebuthiuron (Spike®)
Glyphosate (Razor® Pro)	Triclopyr (Tahoe®)
Hexazinone (Velpar®)	2,4-D
Imazapyr (Arsenal®)	2,4-DP

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies) a COMPATIBILITY TEST is recommended prior to actual tank mixing.

*Restricted use pesticides limited to certified applicators.

** Using Vengeance® Plus and Patriot® to reduce the development and spread of resistant biotypes and problem weeds such as Kochia and Russian thistle: some commonly resistant weeds typically require multiple spray applications to obtain adequate control. To reduce the number of applications required, applicators can utilize a mixture of Patriot® and Vengeance® Plus. Apply as a tank mix for post-emergence as follows: Add .5 to 1 ounce of Patriot® into 1.6 to 4.8 pints of Vengeance® Plus per acre. A CPDA-certified nonionic or silicone surfactant may be used for wetting, penetration or drift control.

NOTE: All intended tank mix combinations should be used only in recommended areas on the same broadleaf weed species found on both labels. For application methods and other use specifications, use the most restricted limitations from labeling of both products.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter Vengeance® Plus. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS", AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS Logo® is a Registered Trademark of Wilbur-Ellis Company. Vengeance® is a registered Trademark of Bayer CropScience. Razor®, The Optical Advantage®, and Patriot® are Registered Trademarks of Nufarm Americas, Inc. Diablo™, Spyder™ and Tahoe® are Trademarks of Nufarm Americas, Inc. Arsenal® is a Registered Trademark of BASF. Princep® is a Registered Trademark of Syngenta Crop Protection. Tordon® and Spike® are Registered Trademarks of Dow AgroSciences LLC. Hyvar®, Karmex®, Krenite®, Telar® and Velpar® are Registered Trademarks of E. I. DuPont de Nemours & Co., Inc. Asulox® is a Registered Trademark of Rhone-Poulenc Chemical Company.

**For Chemical Spill, Leak, Fire or Exposure,
Call CHEMTREC (800) 424-9300**

F-011309-1

Distributed by:

WILBUR-ELLIS COMPANY

PO BOX 16458 • FRESNO CA 93755